

FIELD INVESTIGATION OF *BITHYNIA FUNICULATA*, INTERMEDIATE HOST OF *OPISTHORCHIS VIVERRINI* IN NORTHERN THAILAND

Radchadawan Ngern-klun¹, Kabkaew L Sukontason¹, Smarn Tesana², Duanghatai Sripakdee¹, Kim N Irvine³ and Kom Sukontason¹

¹Department of Parasitology, Faculty of Medicine, Chiang Mai University, Chiang Mai; ²Department of Parasitology, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand; ³Department of Geography and Planning, Buffalo State, State University of New York, Buffalo, NY, USA

Abstract: A survey of freshwater snail, *Bithynia funiculata*, was conducted in four locations, Doi Saket, Mueang, Saraphi and Mae Rim Districts of Chiang Mai Province, northern Thailand, between June and October, during the rainy season of 2004. A total of 2,240 snails was collected and classified into 7 families and 15 genera; of which 352 *B. funiculata* were obtained. *B. funiculata* was found most abundant in July and September. The infection rate of trematode cercariae in *B. funiculata* was 9.6% (19/352), while that of pleurolophocercous was 0.3% (1/352). Virgulate cercaria was the most common type, followed by lophocercous, monostome and pleurolophocercous. *B. funiculata* prefers habitats with clear water, which was less than 30 cm depth, temperatures between 24.48 and 31.78°C, dissolved oxygen 2.03-7.66 mg/l, saturated dissolved oxygen 26.70-95.00%, conductivity 0.000-0.2642 mS/cm, turbidity 16.00-288.00 NTU and pH 6.58-7.56. Geographic Information System (GIS) analysis using loose soil mixed with clay revealed that the breeding grounds of this snail species was in the paddy fields and village environs of the Ping, Kuang and Fang river basins.

INTRODUCTION

The liver fluke, *Opisthorchis viverrini*, is a medically important parasite infecting humans in many countries. It has been estimated by the World Health Organization (1995) that infection caused by this parasite affects approximately 9 million people worldwide. In South-east Asia, endemic areas include Thailand, Cambodia and Lao PDR (Sithithaworn and Haswell-Elkins, 2003). *O. viverrini* has a marked regional variation specifically in Thailand, with a high prevalence occurring in the

northeast and northern regions, where raw and/or undercooked fish is frequently consumed (Sadun, 1995). Epidemiological surveys during 1981-1991 indicated an increasing infection rate of *O. viverrini* in the north and central regions of the country, whereas, rates decreased in the northeastern part (Jongsuksuntigul and Imsomboon, 1998). A recent national survey in 2001 revealed that such infection in the northern region was top ranked, with a prevalence of 19.3% (Jongsuksuntigul, 2002).

The life cycle of *O. viverrini* is governed by the existence of freshwater snails as the first intermediate host and freshwater fish as the second. In Thailand, three taxa of *Bithynia* snails act as the first intermediate host; but, the species of *Bithynia* prevails differently de-

Correspondence: Dr Kom Sukontason, Department of Parasitology, Faculty of Medicine, Chiang Mai University, Chiang Mai 50200, Thailand.
Tel: +66 53 945342; Fax: +66 53 217144
E-mail: ksukonta@mail.med.cmu.ac.th

pending upon geographical habitats; *Bithynia funiculata* in the north, *Bithynia siamensis goniomphalos* in the northeast and *Bithynia siamensis siamensis* in the central part (Brandt, 1974). The natural infection of *O. viverrini* surveyed in these snails varied, ranging from 0.05% to 1.6% (Upatham and Sukhapanth, 1980; Brockelman *et al*, 1986; Adam *et al*, 1993). Using either laboratory or field collected snails, *B. funiculata* was significantly more susceptible to *O. viverrini* infection than both *B. goniomphalos* and *B. siamensis* (Chanawong and Waikagul, 1991).

Ecological environment has been recorded as having influence on the fluctuation of the snail as an intermediate host in an endemic area (eg, Upatham and Sukhapanth, 1981; Brockelman *et al*, 1986; Wullschlegler and Jokela, 1999; Kock *et al*, 2004; Rondelaud *et al*, 2004). Nevertheless, information pertaining to water quality as well as the physical environment of the breeding grounds of *B. funiculata* in Chiang Mai, northern Thailand, is very limited. In this study, we surveyed the prevalence of *B. funiculata* and the infection rate of trematode cercariae in this snail species in Chiang Mai during the rainy season of 2004. Other freshwater snails inhabiting the same location with *B. funiculata* were documented. We also investigated the ecological factors, water quality and physical environment, that may entail suitable habitats for *B. funiculata*, while using the application of Geographical Information System (GIS) technology to incorporate snail survey data and environmental factors to predict the distribution of *B. funiculata* in the study areas of Chiang Mai.

MATERIAL AND METHODS

Study areas and land use

Four study areas (Fig 1) were chosen around the downtown area of Chiang Mai (17-21°N, 98-99°E) to investigate the *B. funiculata*

breeding sites. Area 1 was located at Ban Thawang (Mae Rim District). Areas 2, 3 and 4 were located at Ban Nhongpom (Mueang District), Ban Pakui (Doi Saket District) and Ban Si Boonreang (Saraphi District), respectively (Fig 1A). Descriptive information of such areas is given in Table 1. The coordinates for each site were recorded using the Global Positioning System (GPS) Garmin® GPS 12 Personal Navigator® (Garmin International, Kansas, USA). Rainfall data were obtained from the Meteorological Center, Chiang Mai, Thailand. The depth of water, characteristics of mud, sand and soil substrata of each site was recorded.

Snail sampling method

Routine sampling of snails was conducted monthly from June to October 2004, which represented the rainy season. Snail collection was performed twice a month. The snail sampling method employed was dependent upon the ecology of each study site. Hand scoop sampling was performed in the area having aquatic plants by using the standard wire-mesh scoop with a mesh size of 2 mm. At each sampling location, ten scoops were made along just beneath the mud surface in a set pattern (4x4 m). Thereafter, the mud was rinsed off the scoops and the snails were collected one by one. In areas without aquatic plants, the snails were picked up manually by two people for 10 minutes at each sample site.

After collection, the snails were placed in a plastic bag and transferred to the Department of Parasitology, Faculty of Medicine, Chiang Mai University. All specimens were identified by the methods of Brandt (1974) and Upatham *et al* (1983). Only *B. funiculata* was measured for its size and examined for trematode cercariae shedding.

Cercarial shedding

B. funiculata collected from each site was placed separately in a test tube containing

Table 1
Description of study areas where snail collections were conducted from June to October 2004 in Chiang Mai, northern Thailand.

Area	District	Village	Sub-sites	Soil type	Water supply	Altitude ^a	Georeference coordinates
I	Mae Rim	Ban Thawang	S1P1 (rice paddy)	Series-5 (loose soil + clay)	Suthep Mountain and Ping river	319	49.70 E; 20.91 N
			S1P2 (rice paddy)				
			S1C (irrigation canal)				
II	Mueang	Nong Pom	S2P (rice paddy)	Series-7 (sandy soil + clay)	Suthep Mountain and Ping river	317	49.93 E; 20.85 N
			S2C (irrigation canal)				
			S3P (rice paddy)				
III	Doi Saket	Pakui	S3C (irrigation canal)	Series-59 (clay)	Mae Kuang Udom Thara Dam	315	50.59 E; 20.80 N
			S4P (rice paddy)				
IV	Saraphi	Si Boonreang	S4P (rice paddy)	Series-5 (loose soil + clay)	Mae Kuang Udom Thara Dam	314	50.54 E; 20.72 N

^aMeters above sea level

distilled water. Such tubes were exposed to the illumination of a 25-W light source at a distance of 30 cm for 2 hours at room temperature. Subsequently, the water from each tube was poured into a Petri dish and examined for cercarial shedding under a dissecting microscope. Identification of cercariae was performed by pipetting a few drops of water onto a glass slide, covered with a cover slip, and then examining under a stereomicroscope, as described by Frandsen and Christensen (1984).

Water quality recording

The water quality of each sample site during every snail collection was monitored and recorded with Hydrolab Datasonde® 4a (Hydrolab corporation, Texas, USA). The parameters recorded were temperature, dissolved oxygen (both concentration and percent of saturation), conductivity, turbidity and pH.

GIS database

Base topographical maps of the four locations in this study were digitalized, input and overlaid, with the data gained from each site entered into a GIS database using ArcView 3.3 software. Other thematic maps [soil map, water supply, land-use classes of paddy field (Fig 1C)] were also incorporated. All digital data were displayed in the UTM coordinate system and georeferenced using the Indian 1975 Datum and Everest 1830 (1975 Definition) Ellipsoid Provincial border boundary.

RESULTS

A total count of 2,240 snail specimens was collected from four localities in Chiang Mai Province during the rainy season from June to October 2004. Seven families comprised of fifteen genera were obtained (Table 2). Within the family Bithyniidae, *Hydrobioides nassa* ranked the most prevalent (26.2%), followed by *Wattebledia siamensis* (17.6%), *B. funiculata* (15.7%) and *Bithynia (Gabbia)*

Table 2

Collected freshwater snail from four study areas around downtown Chiang Mai, from June to October 2004.

Family	Snail collected		
	Species	N	Percentage
Ampullariidae	<i>Pomacea</i> spp	75	3.3
Bithyniidae	<i>Bithynia (Digoniostoma) funiculata</i>	352	15.7
	<i>Bithynia (Gabbia) wykoffi</i>	59	2.6
	<i>Hydrobioides nassa</i>	586	26.2
	<i>Wattebledia siamensis</i>	394	17.6
Buccinidae	<i>Clea (Anentome) helena</i>	35	1.6
Lymnaeidae	<i>Austropeplea ollula</i>	95	4.2
Planorbidae	<i>Camptoceras jiraponi</i>	3	0.1
	<i>Indoplanorbis exustus</i>	11	0.5
Thiaridae	<i>Melanooides tuberculata</i>	301	13.4
	<i>Tarebia granifera</i>	70	3.1
	<i>Thiara scabra</i>	2	0.1
Viviparidae	<i>Eyriesia</i> sp	1	0.0
	<i>Filopaludina</i> spp	135	6.0
	<i>Idiopoma</i> spp	44	2.0
Unidentified		77	3.4
Total		2,240	100.0

wykoffi (2.6%), consecutively.

Of the 352 *B. funiculata* sampled, most of the snails were collected in July (n = 115) and September (n = 102) when rainfall was highest (Table 3). However, there was no correlation between the number of snails and quantity of rainfall (p = 0.391). *B. funiculata* snails were present in 3 Areas of 1, 3 and 4 (Table 3). In the paddy fields of Areas 1 and 4, where soil characteristics were graded as loose soil and clay, *B. funiculata* was found in all five months of the study period. In Area 3, which had a clay type of soil (series 59), *B. funiculata* could be collected only from August to October. No *B. funiculata* was found in the irrigation canals of Area 1 and 3. Likewise, snails could not be detected in the paddy fields of Area 3, which had sandy and clay soil types.

The infection rate of trematode cercariae in *B. funiculata* was 5.4% (19/352) (Table 4). Large *B. funiculata* ($\phi > 7.0$ mm) had an infec-

tion rate of 4.8%, while 0.5% was found in smaller ones ($\phi \leq 7.0$ mm). However, the infection rate in large and small snails was not statistically significant (p = 0.071). Of the nineteen infected *B. funiculata*, virgulate cercaria was the most common type (63.2%), followed by lophocercous (21.0%), monostome (10.5%) and pleurolophocercous cercaria (5.3%) (Table 5). The pleurolophocercous cercaria was probably the cercaria of *O. viverrini*. No mixed infection of trematode cercariae was detected in the same snail specimen. The number of pleurolophocercous cercaria released from a single snail was 286. Morphological features and measurement of these four types of cercaria are shown in Table 6. Monostome cercaria had the biggest body and a distinctive three-eyed spot. Lophocercous and pleurolophocercous cercaria had the same body length; however, the former was broader than the latter (119.26 ± 18.90 vs 57.87 ± 11.76 μ m).

Table 3

Number of *B. funiculata* collected from four area of Chiang Mai from June to October 2004.

Month	Area 1			Area 2		Area 3		Area 4	Total	Rainfall ^a (mm)
	S1P1	S1P2	S1C	S2P	S2C	S3P	S3C	S4P		
June	9	6	0	0	0	0	0	15	30	178.8
July	60	45	0	0	0	0	0	10	115	218.0
August	6	16	2	0	0	7	0	1	32	115.7
September	20	8	12	0	0	22	0	40	102	317.4
October	14	5	0	0	0	8	0	46	73	38.3
Total		203		0		37		112	352	

^a There was no correlation between the number of snails collected and rainfall ($p = 0.391$; $r = 0.50$, Spearman rank correlation coefficients)

Table 4

Infection rate of trematode cercariae found in field collected *B. funiculata* ranked by size.

Length of snails (mm)	Total snails		No. of <i>B. funiculata</i>		Infection rate (%)
	No.	%	Uninfected	Infected	
≤ 7.00	108	30.7	106	2	0.6
> 7.00	244	69.3	227	17	4.8
Total	352	100.0	333	19	5.4
p					0.071 ^a

^a No significant difference ($p > 0.05$; chi-square tests)

Table 5

Type of cercariae recovered from *B. funiculata* collected in study areas around downtown Chiang Mai, from June to October 2004.

Type of cercariae	Number of infected snails (%)	Rate of infection (%)	No. shed cercariae/snail (range) (%)
Virgulate	12 (63.2)	3.4	452 (1-1,241)
Lophocercous	4 (21.0)	1.1	37 (7-197)
Monostome	2 (10.5)	0.6	56 (28-84)
Pleurolophocercous	1 (5.3)	0.3	286
Total	19 (100)	5.4	

Table 7 displays information pertaining to water quality recorded from areas where *B. funiculata* was collected. It was found that *B. funiculata* was present in water at temperatures between 24.48 and 31.78°C, DO rang-

ing from 2.03-7.66 mg/l, DO% of 26.70-95.00, conductivity between 0.000 and 0.264 mS/cm, turbidity between 16.00 and 288.00 NTU, and pH of 6.58-7.56. Moreover, *B. funiculata* preferred water less than 30 cm depth. Areas

Table 6
Characteristics of trematode cercaria recovered from *B. funiculata*.

Type of cercariae	Body (m) (mean ± SD)		Tail (μ) (mean ± SD)		Special characteristics
	Length	Width	Length	Width	
Virgulate	108.07± 11.89	59.59± 12.13	55.8± 11.82	10.79± 3.00	Oral and ventral suckers were large; the ventral were smaller than the oral. Single stylet displayed obvious anteriority. Tail was relatively short with no finfold.
Lophocercous	178.57± 35.93	119.26± 18.90	417.1± 38.12	30.26± 4.5	Body was oval. Oral and ventral suckers, eye spot and transverse array finfold were apparent.
Monostome	323.78± 54.62	159.34± 23.28	499.87± 102.24	30.61	Distinct three-eyed spot and well-developed oral sucker. Tail had no finfold.
Pleurophocercous	183.67± 23.55	57.87± 11.76	417.1± 38.12	30.26± 4.50	Body was cylindrical. Lateral finfold was prominent.

Table 7
Water quality recorded from sites where *B. funiculata* was collected.

<i>B. funiculata</i>	Median (range)				
	Temperature (°C)	Dissolved oxygen (%)	Dissolved oxygen (mg/l)	Conductivity (mS/cm)	Turbidity (NTU)
Presence	26.51 (24.48-31.78)	47.45 (26.70-95.00)	3.79 (2.03-7.66)	0.147 (0.000-0.264)	68.55 (16.00-288.00)
Absence	27.86 (24.71-30.42)	59.90 (18.30-83.70)	4.71 (1.42-10.44)	0.175 (0.000-0.366)	31.25 (13.00-92.00)
p	0.89	0.126	0.154	0.196	0.001 ^a

^a Significant difference; Mann-Whitney U test (p < 0.05)




Fig 1—Location of four collection areas in Chiang Mai, northern Thailand, where snails were collected from June to October, 2004. (1A) 1 = Area 1, Ban Thawang, Mae Rim District; 2 = Area 2, Ban Nhongpom, Mueang District; 3 = Area 3, Ban Pakui, Doi Saket District; 4 = Area 4, Ban Si Boonreaung, Saraphi District. (1B) Thematic map showing soil type series 5 and 59. (1C) Thematic map showing paddy field areas. (1D) Predicted area of Chiang Mai where *B. funiculata* is present.

where *B. funiculata* snails were found had significantly higher turbidity and lower pH than those without *B. funiculata* ($p = 0.001$ and 0.012 , respectively).

Using the Arcview 3.3 program to incorporate the snail survey data with soil type (series 5 and 59: Fig 1B), land used (paddy field) and villages of Chiang Mai, the probable breeding grounds of *B. funiculata* were predicted as the Ping, Kuang and Fang river basins and encompassed 165 villages in Chiang Mai. Nevertheless, the expected areas were mainly positioned in the Ping and Kuang river basins (Fig 1D).

DISCUSSION

Many freshwater snail species were present in this study, including *B. funiculata*, the first intermediate host of an important liver fluke in Thailand. The population of *B. funiculata* collected in the months of heavy rain was high, but there was no statistically significant correlation between snail population and rainfall. This phenomenon was unlike the collections of *Bithynia siamensis siamensis* or *Indoplanorbis exustus* investigated in Bangkok, and central Thailand (Upatham and Suthapanth, 1980, 1981). It was interesting to note that heavy rains did not affect the collection of *B. funiculata* in this study. The consistent water depth maintained by

the irrigation systems of the paddy fields and the presence of more snails in soil substrate than at the surface level of water may explain the lack of correlation. *B. funiculata* could be

found in all months of the rainy season in paddy fields that had loose soil and clay. However, in sub-site S3P of Area 3, which was a paddy field with clay, *B. funiculata* was found at depths of 30 cm in clear water after rice had been planted.

Regarding the natural parasitic infection of *B. funiculata*, snails liberated four types of trematode cercariae: virgulate, lophocercous, monostome and pleurolophocercous cercariae. Ito *et al* (1962) and Wykoff *et al* (1965), classified *O. viverrini* as a type of pleurolophocercous cercariae. With respect to *B. goniomphalos*, the types of cercariae liberated were lophocercous, monostome, furcocercous, xiphidiocercariae (Adam *et al*, 1993), echinostomes, intestinal flukes, xyphidia, furcocercous, *O. viverrini* (Sriaroon *et al*, 2005) or amphistome (Wiwanitkit *et al*, 2002). With respect to *B. siamensis*, the type of cercarial emitted were furcocercous and pleurolophocercous (*O. viverrini*) (Upatham and Sukhapanth, 1980).

The infection rate of pleurolophocercous (*O. viverrini*) cercaria in naturally infected *B. funiculata*, as observed in this study, was relatively low (0.28%) (Table 4), and lower than those of *B. siamensis* in central Thailand (1.6%) (Upatham and Sukhapanth, 1980), *B. goniomphalos* in Lao PDR (0.5-0.9%) (Ditrich *et al*, 1990; Giboda *et al*, 1991). However, a survey of *O. viverrini* cercaria from field collected *B. goniomphalos* in northeast Thailand exhibited no infection, despite the high prevalence of *O. viverrini* infection in the studied area (Nithiuthai *et al*, 2002). This was consistent the 0.11% and 0.05% minimal infection rate of *O. viverrini* in *B. goniomphalos* reported by Brockelman *et al* (1986) and Adam *et al* (1993), respectively.

Larger (>7.0 mm) field-collected *B. funiculata* yielded a higher infection rate of *O. viverrini* than smaller ones (Table 4). Similarly, Upatham and Sukhapanth (1980) observed more infections by this parasite in larger *B.*

siamensis (>7.0 mm). Interestingly, a study by Chanawong and Waikagul (1991) found that the immature laboratory bred *Bithynia* (2-4 mm long) was more susceptible to *O. viverrini* infection than mature field-collected specimens (6-10 mm long).

Field investigations in this study indicated that *B. funiculata* inhabit rice paddy fields with clear water of no more than 30 cm in depth. Most *B. funiculata* were sampled from the muddy substrate of rice paddies (S1P, S2P, S4P), which is consistent with previous reports (TROPED Technical Group, 1986; Chitramvong, 1992). *B. funiculata* was generally observed, as in the collection sites S1P, S2P and S4P, sliding over mud substrate or attached to the bottom of rice, aquatic vegetation and algae. The swampy area of the Bhumibol reservoir (Tak Province, northern Thailand) has also been reported as a habitat of *B. funiculata* infected with *O. viverrini* (Rojanapremsuk *et al*, 1988). Other members belonging to the family Bithyniidae, including *Hydrobioides nassa* and *Wattebledia siamensis*, were found in similar habitats. In general, the ecological niches of *B. funiculata* in this study were similar to those of *B. goniomphalos* investigated in northeast Thailand, in which snails inhabit static water at a depth of no more than 30 cm along the edge of a water reservoir. They were found sliding on aquatic plants growing in red-yellow podzolic soils (Papasarathorn *et al*, 1990). Nevertheless, *B. goniomphalos* have been found in other habitats (Brockelman *et al*, 1986), including mixed sand-mud substrate at pond's edge, where the water is relatively deep and devoid of algae (Chitramvong, 1992).

Determination of water quality in each collection site revealed that the turbidity and pH of the water were the principle parameters which determined the presence and absence of *B. funiculata*. *B. funiculata* preferred water with high turbidity and a pH of 6.58-7.56. This

pH value was broader than the 6.90-7.10 pH value earlier reported for bithyniid snails (TROPMED Technical Group, 1986), or the 6.6-7.2 pH of specific *B. goniomphalos* (Parasarathorn *et al*, 1990). *B. goniomphalos* were also presented in water with high turbidity (Parasarathorn *et al*, 1990).

GIS technology has been applied to investigate and understand the ecology of the intermediate host of medically and/or veterinarily important parasites, such as the snail intermediate hosts of the liver fluke, *Fasciola hepatica* (Zukowski *et al*, 1991, 1993; Yilma and Malone, 1998) and *Fasciola gigantica* (Yilma and Malone, 1998); the blood fluke, *Schistosoma mansoni* (Kristensen *et al*, 2001), *Schistosoma haematobium* (Lwambo *et al*, 1999), and *Schistosoma japonicum* (Maszle *et al*, 1998; Yang *et al*, 2005; Nihei *et al*, 2006); and rumen fluke, *Calicophoron daubneyi* (Cringoli *et al*, 2004). Our report, presented herein, is the first recorded GIS-facilitated monitoring of the snail intermediate host of *O. viverrini* in Chiang Mai. Analysis indicates that the habitat of *B. funiculata* in Chiang Mai is characterized by paddy fields with loose soil mixed with clay. Based on our data, we hypothesize that the *B. funiculata* population is distributed in the central part of Chiang Mai Province. The collection of additional data in similar habitats is an ongoing project and is needed to confirm this hypothesis.

ACKNOWLEDGEMENTS

We are grateful to Chira Prangki and Kittisak Chaokasad of the Department of Geography, Faculty of Social Science, Chiang Mai University, for kindly providing the database of thematic maps. We also thank Amy M Krueger of the Department of Geography and Planning, State University of New York, for her kind assistance. We acknowledge the Faculty of Medicine and Chiang Mai University for

meeting the publication cost of this paper.

REFERENCES

- Adam R, Arnold H, Pipitgool V, Sithithaworn P, Hinz E, Storch V. Studies on lophocercous cercariae from *Bithynia siamensis goniomphalos* (Prosobranchia: Bithyniidae). *Southeast Asian J Trop Med Public Health* 1993; 24: 697-700.
- Brandt RAM. The non-marine aquatic mollusca of Thailand. *Arch Molluskenkd* 1974; 105: 1-423.
- Brockelman WY, Upatham ES, Viyanant V, Ardsungnoen S, Chantanawat R. Field studies on the transmission of the human liver fluke, *Opisthorchis viverrini*, in northeast Thailand: population changes of the snail intermediate host. *Int J Parasitol* 1986; 16: 545-52.
- Chanawong A, Waikagul J. Laboratory studies on host-parasite relationship of *Bithynia* snails and the liver fluke, *Opisthorchis viverrini*. *Southeast Asian J Trop Med Public Health* 1991; 22: 235-9.
- Chitramvong YP. The Bithyniidae (Gastropoda: Prosobranchia) of Thailand: comparative external morphology. *Malacol Rev* 1992; 25: 21-38.
- Cringoli G, Taddei R, Rinaldi L, *et al*. Use of remote sensing and geographical information systems to identify environmental features that influence the distribution of paramphistomosis in sheep from the southern Italian Apennines. *Vet Parasitol* 2004; 122: 15-26.
- Ditrich O, Scholz T, Giboda M. Occurrence of some medically important flukes (Trematoda: Opisthorchiidae and Heterophyidae) in Nam Ngum water reservoir, Laos. *Southeast Asian J Trop Med Public Health* 1990; 21: 482-8.
- Frandsen F, Christensen NO. An introductory guide to the identification of cercariae from African freshwater snails with special references to cercariae of trematode species of medical and veterinary importance. *Acta Trop* 1984; 41: 181-202.
- Giboda M, Ditrich O, Scholz T, Viengsay T, Bouaphanh S. Current status of food-borne parasitic zoonoses in Laos. *Southeast Asian J*

- Trop Med Public Health* 1991; 22 (suppl): 56-61.
- Ito J, Papasarathorn T, Tongkoom B. Studies on cercariae from freshwater snails in Thailand. *Jpn J Med Sci Biol* 1962; 15: 249-70.
- Jongsuksuntigul P. Parasitic diseases in northeast Thailand. In: Seminar in Parasitic Diseases in Northeast Thailand, Khon Kaen, Thailand, 22-23 November 2002: 3-18.
- Jongsuksuntigul P, Imsomboon T. Epidemiology of opisthorchiasis and national control program in Thailand. *Southeast Asian J Trop Med Public Health* 1998; 29: 327-32.
- Kock KN, Wolmarans CT, Bornman M. Distribution and habitats of *Biomphalaria pfeifferi*, snail intermediate host of *Schistosoma mansoni*, in South Africa. *Water SA* 2004; 30: 29-35.
- Kristensen TK, Malone JB, McCarroll JC. Use of satellite remote sensing and geographical information systems to model the distribution and abundance of snail intermediate hosts in Africa: a preliminary model for *Biomphalaria pfeifferi* in Ethiopia. *Acta Trop* 2001; 79: 73-8.
- Lwambo NJS, Siza JE, Brooker S, Bundy DAP, Guyatt H. Patterns of concurrent hookworm infection and schistosomiasis in schoolchildren in Tanzania. *Trans R Soc Trop Med Hyg* 1999; 93: 497-502.
- Maszle DR, Whitehead PG, Johnson RC, Spear RC. Hydrological studies of schistosomiasis transport in Sichuan Province, China. *Sci Total Environ* 1998; 216: 193-203.
- Nihei N, Kajihara N, Kirinoki M, *et al.* Establishment of a GIS monitoring system for schistosomiasis japonica in Kofu, Japan. *Ann Trop Med Parasitol* 2006; 100: 143-53.
- Nithiuthai S, Wiwanitkit V, Suwansaksri J, Chaengphukeaw P. A survey of trematode cercariae in *Bithynia goniomphalos* in northeast Thailand. *Southeast Asian J Trop Med Public Health* 2002; 33 (suppl 3): 106-9.
- Papasarathorn T, Sumethanurugkul P, Viboolyavatana J, Temcharoen P, Tongkoom B. Studies of life history and ecological requirements of *Bithynia (Digoniostoma) siamensis goniomphalos*, the snail intermediate host of *Opisthorchis viverrini*. *J Parasitol Trop Med Assoc Thai* 1990; 3: 21-7.
- Rojanapremsuk J, Temcharoen P, Viboolyavatana J, Pandii W. Study on parasitic infections of the people around the Bhumibol and the Sirikit water resource development projects, northern Thailand. *J Parasitol Trop Med Assoc Thai* 1988; 11: 62-70.
- Rondelaud D, Vignoles P, Dreyfuss G. *Fasciola hepatica*: the developmental patterns of redial generations in naturally infected *Galba truncatula*. *Parasitol Res* 2004; 94: 183-7.
- Sadun EH. Studies on *Opisthorchis viverrini* in Thailand. *Am J Hyg* 1955; 62: 81-115.
- Sithithaworn P, Haswell-Elkins M. Epidemiology of *Opisthorchis viverrini*. *Acta Trop* 2003; 88: 187-94.
- Sri-aroon P, Butraporn P, Limsomboon J, Kerdpuech Y, Kaewpoolsri M, Kiatsiri S. Freshwater mollusks of medical importance in Kalasin Province, northeast Thailand. *Southeast Asian J Trop Med Public Health* 2005; 36: 653-7.
- TROPED Technical Group. Snails of medical importance in Southeast Asia. *Southeast Asian J Trop Med Public Health* 1986; 17: 282-322.
- Upatham ES, Sornmani S, Kitikoon V, Lohachit C, Burch JB. Identification key for the fresh- and brackish-water snails of Thailand. *Malacol Rev* 1983; 16: 107-32.
- Upatham ES, Sukhapanth N. Field studies on the bionomics of *Bithynia siamensis siamensis* and the transmission of *Opisthorchis viverrini* in Bangna, Bangkok, Thailand. *Southeast Asian J Trop Med Public Health* 1980; 11: 355-8.
- Upatham ES, Sukhapanth N. Field studies on the bionomics of freshwater snails of medical importance in Bangna, Thailand. *Southeast Asian J Trop Med Public Health* 1981; 12: 204-8.
- Wiwanitkit V, Suwansaksri J, Nithiutha S. Prevalence of larval helminthes in fresh water snails, *Bithynia goniomphalos* of Nong Rua, Khon Kaen, Thailand. *Songkla Med J* 2002; 20: 95-8.

- World Health Organization. Control of foodborne trematode infections. Report of a WHO study group. *WHO Techn Rep Ser* 1995; 849: 1-157.
- Wullschleger E, Jokela J. Does habitat-specific variation in trematode infection risks influence habitat distribution of two closely related freshwater snails? *Oecologia* 1999; 121: 32-8.
- Wykoff DE, Harinasuta C, Juttijudata P, Winn MM. *Opisthorchis viverrini* in Thailand – The life cycle and comparison with *O. felineus*. *J Parasitol* 1965; 51: 207-14.
- Yang GJ, Vounatsou P, Zhou XN, Tanner M, Utzinger J. A potential impact of climate change and water resource development on the transmission of *Schistosoma japonicum* in China. *Parassitologia* 2005; 47: 127-34.
- Yilma JM, Malone JB. A geographical information system forecast model for strategic control of fasciolosis in Ethiopia. *Vet Parasitol* 1998; 78: 103-27.
- Zukowski SH, Hill JM, Jones FW, Malone JB. Development and validation of a soil-based geographic information system model of habitat of *Fossaria bulimoides*, a snail intermediate host of *Fasciola hepatica*. *Prev Vet Med* 1991; 11: 221-7.
- Zukowski SH, Wilkerson GW, Malone JB Jr. Fasciolosis in cattle in Louisiana. II. Development of a system to use soil maps in a geographic information system to estimate diseases risk of Louisiana coastal marsh rangeland. *Vet Parasitol* 1993; 47: 51-65.